

The current state of COVID-19 in Colorado

9/11/2020

Prepared by the Colorado COVID-19 Modeling Group

Colorado School of Public Health: Andrea Buchwald, Elizabeth Carlton, Debashis Ghosh, Jonathan Samet, Laura Timm, Emily Wu; University of Colorado School of Medicine: Kathryn Colborn; University of Colorado-Boulder Department of Applied Mathematics: David Bortz; University of Colorado-Denver: jimi adams; Colorado State University: Jude Bayham

For Contact: Jon.Samet@CUAnschutz.edu

Summary

- COVID-19 hospitalizations have been decreasing since mid-July. This decline has slowed and hospitalizations appear to be at a plateau.
- The estimated effective reproductive number has increased since last week following an upward trend over recent weeks. We estimate the current effective reproductive number is between 0.93 and 1.14. Due to lags between infection and hospitalizations, this reflects transmission occurring through late-August (approximately 8/25).
- The model-estimated number of infectious individuals in Colorado remains relatively low. The model-estimated rate of new infections as well as reported cases had been declining since mid-July and now appears to be leveling off.
- Our current estimate of social distancing is between 65% for the week of 8/17 - 8/25, and 72% during the three-week period 8/03 - 8/25. If we remain at 65%, we will begin to see gradual growth in cases and/or hospital demand.
- Colorado colleges and universities continue to report increases in new cases. We are monitoring this closely.

Current state of COVID-19 in Colorado

We used our age-structured SEIR model and COVID-19 hospital census data to characterize the current status of the COVID-19 epidemic in Colorado and the collective impact of efforts to date to reduce the spread of the SARS-CoV-2 virus. Modeling methods are detailed [here](#), (see documentation tab). These estimates are based on hospitalization data through 09/07/2020.

To characterize the current state of COVID-19 in Colorado, we focus on three components:

1. **The effective reproductive number.** This is an estimate of the recent growth rate of infections. If the reproductive number is above 1, the number of infections is growing, with higher values indicating faster growth. If the reproductive number is below 1, cases are declining.
2. **The cumulative and current number of infections in the population.** The current number of people who are infectious at a given time provides a measure of the risk of infection to the population at large. When the number of infectious individuals is high, there is a greater chance that an interaction between people may lead to a new infection. The cumulative number of infections to date provides an estimate of the size of the remaining susceptible population, the population that remains at risk of infection, assuming there is durable immunity.

3. **The distribution of reported infections and hospitalizations by age, race and ethnicity.** The impacts of COVID-19 have not been distributed evenly across the population. By examining cases and hospitalizations by age, race and ethnicity over time, we can identify high-risk populations.

These estimates are based on currently available data and our current understanding of SARS-CoV-2 transmission. We note that the science on COVID-19 and the pandemic are evolving rapidly, and we regularly update our model assumptions as new information becomes available.

Current COVID-19 hospitalizations and model fit

Figure 1 shows COVID-19 hospitalizations (black bars) and the green line shows the current model fit to the data. Table A1 provides values for model parameters including social distancing estimated by the model from mid-March through the present. The social distancing parameters describe reductions in the contact rate, a measure of how rapidly infections spread from person to person. Our most recent estimate of social distancing, for the period 8/17 to 8/25, is 65%. We note that due to the approximately 13-day lag between infection and hospitalization, we are currently only able to estimate social distancing through 8/25.




Figure 1. Current model fit (green line) to count of hospitalized COVID-19 cases (black lines) using the age-structured SEIR model. Hospitalized COVID-19 cases are from CDPHE reported COVID-19 hospitalizations and EMResource (EMR) hospital census data provided by CDPHE.

The effective reproductive number

The estimated effective reproductive number is shown in Table 1 and Figure 2. Table 1 provides estimates we generated using two different methods. We also provide values from two external groups that use different data and methods (RT-Live uses SARS-Cov-2 case data and covid-19-projections.com uses mortality data). Trends in these external estimates reflect our estimates.

Table 1. Current and prior estimates of the effective reproductive number (R_e) in Colorado.

	Current Estimate (09/07)	Estimate one week ago (8/31)	Estimate two weeks ago (8/24)
Estimate of R_e , approach 1*	1.08	0.87	0.79
Estimate of R_e , approach 2*	1.14	0.80	0.76
Estimate from RT-Live	0.99	0.93	0.90
covid-19-projections.com	0.93	0.99	0.94

*Our estimates are based on hospitalization data through the date listed. Estimates from the external sites are extracted on the day listed. Because of the 13-day lag between infection and hospitalization, on average, our current estimate reflects transmission up to approximately August 25. Approach 1 uses model output to estimate the average number of new cases generated by existing cases, accounting for the latent period and duration of infectiousness. The second method uses the model structure to estimate the dominant eigenvalue for a matrix describing population flows across the model compartments.




Figure 2. The effective reproductive number using approach 1 (top panel) and approach 2 (bottom panel) based on our SEIR model.




Figure 3. The projected trajectory of COVID-19 hospitalizations if Colorado remains on the current estimated trajectory (orange line), the trajectory estimated one week prior (red line), and the trajectory estimated two weeks prior (blue line). Each trajectory is generated assuming social distancing levels remain at the estimated levels: current estimate 65% based on the period 8/17 - 8/25, one week prior estimate 73% based on the period 8/03 - 8/18 (red line), two weeks prior estimate 75% based on the period 8/03 - 8/11 (blue line). Note that the estimation periods overlap as we re-estimate parameters each week and use the past approximately 10 days to estimate the most recent social distancing parameter.

The cumulative and current number of infections in the population

We use our model output to estimate the cumulative number of infections to date and the approximate number of infectious individuals in the population. Given the characteristics of SARS-CoV-2 and of COVID-19, many infections are not detected by surveillance systems – the estimates provided here are intended to provide an approximation of the total number of infections, as well as the proportion detected by the Colorado surveillance system. These estimates are sensitive to our model assumptions, including assumptions about the probability an infected individual will be symptomatic and require hospital care, which we assume varies by age.

We estimate that approximately 294,000 people in Colorado, or 5.0% of the population have been infected to date.

We estimate that there are approximately 5,200 infectious individuals in Colorado at present: approximately 1 in every 1,100 Coloradans.

We estimate that approximately 48% of infections were detected in the past two weeks, including both asymptomatic and symptomatic infections. This estimate is generated by comparing daily reported SARS-CoV-2 cases in Colorado to model-estimates of the number of incident infections.




Figure 4. Estimated daily number of new (incident) SARS-CoV-2 infections based on the SEIR model (dotted line) and reported cases (solid line) over time shown in the top panel. Lower panel shows the estimated proportion of SARS-COV2 infections that are being captured by Colorado state surveillance systems, over time. The proportion detected is estimated by dividing the total number of new cases captured by state surveillance systems by the model-estimated number of new infections each day. The number of cases captured by state surveillance systems is the number of cases reported by CDPHE, using the onset date of symptoms (if onset date is not available, onset date is imputed by CDPHE using a proxy distribution of recent onset dates). Data are shown through 8/25, to account for typical lags between symptom onset and case report.

The distribution of reported infections and hospitalizations by age, race and ethnicity

Figure 5 shows the weekly number of reported new SARS-CoV-2 infections by age group. Recent reports of new cases are highest for those aged 20 - 39. The average proportion of COVID-19 cases in people under age 40 over the last two weeks is approximately 58%.

Figure 6 shows the weekly number of new COVID-19 hospitalizations by age group. Recent new hospitalizations are highest for those age 65 and over. The average proportion of COVID-19 hospitalizations in people under age 40 over the last two weeks is approximately 20%.





Figure 5. Distribution of weekly new (incident) reported SARS-CoV-2 cases by age group (top) and the proportion of all cases among individuals under 40 (bottom). Reported cases are based on CDPHE data

and shown by onset date (if onset date is not available, onset date is imputed by CDPHE using a proxy distribution of recent onset dates). Incident cases per 100,000 were obtained by standardizing weekly reported age-specific case and hospitalization counts to the Colorado population distribution by age, gathered from the Colorado Census 2020 estimates. Data are shown through 8/25/2020, to account for typical lags between symptom onset and case report.





Figure 6. Distribution of weekly new (incident) COVID-19 hospitalizations by age (top) and the proportion of hospitalizations occurring among individuals under 40 (bottom). Hospitalizations per 100,000 were estimated by standardizing weekly reported age-specific hospitalizations based on CDPHE case reports to the Colorado population distribution by age, gathered from the Colorado Census 2020 estimates. Data are shown through 8/25/2020.





Figure 7. Distribution of weekly new (incident) reported SARS-CoV-2 infections (top) and new hospitalizations (bottom) by race and ethnicity in Colorado. Reported cases are based on CDPHE data and shown by onset date (if onset date is not available, onset date is imputed by CDPHE using a proxy distribution of recent onset dates). Cases and hospitalizations per 100,000 were obtained by standardizing weekly reported race-specific case and hospitalization counts to the race/ethnicity distribution of the state of Colorado gathered from the CDPHE COVID-19 Case Summary Dashboard. These standardized estimates combine Asian and Native Hawaiian/Pacific Islander races and exclude Other/Unknown races (which account for 28% of observations over the last two weeks).

What do we project for the coming months?

We used the age-structured SEIR model fit to the parameters described in Table A1 to project the expected number of total hospitalizations and need for critical care beds in the coming months for two sets of scenarios: 1) current and recent trajectories; 2) hypothetical relaxation of social distancing

Current and recent trajectories. Sets of projections are generated assuming we remain on the current trajectory and accounting for uncertainty in our current estimated trajectory (Figure 8). As hospitalization numbers have fluctuated in recent weeks, leading to variation in estimates of social distancing, we estimate current trajectory two ways: first examining the trajectory based on the most recent social distancing parameter estimated for the most recent period, 08/17 – 8/25, and second by using the average social distancing parameter estimated for the last three weeks (the period 08/03 to 08/25). Projections assuming, we remain on our current trajectory or the three-week average trajectory, indicate that we may see a surge in infections but are unlikely to exceed hospital capacity in the next 8 weeks. Projections out past the next 8 weeks are hypothetical and highly uncertain as trajectories change frequently.






Figure 8. Projected daily count of new infections (top), hospital (center), and intensive care (ICU) demand (bottom) in the near-term (left) or long term (right) if we remain on the current trajectory estimated based on the most recent social distancing estimate (purple line, 65% for the period 8/17 – 8/25), and based on the three-week average (blue line, 72% for the period 08/03 – 8/25). Dashed and dotted lines represent a 5% overestimate / underestimate respectively in the social distancing parameter. These scenarios assume mask wearing increases to 90% on 7/16, the date of the state-wide mask order, and case isolation remains at current levels. These projections use model-fit parameters through 8/25 and then switch to the projected parameter values.

Relaxation of social distancing. Projections were also generated to evaluate the potential impact of hypothetical changes to the current trajectory – looking at the potential impact of theoretical reductions in social distancing (Figure 9 and Table 2). For the purpose of these projections, we assume social distancing is relaxed on 9/11. Projections are generated assuming mask wearing increased to 90% on 7/16, the date of the statewide mask order.

Projections show that if social distancing were theoretically relaxed to 55 or 50%, we could see substantial growth in cases in the months ahead, with the sharpest growth at the lowest levels of social distancing. ICU capacity is expected to be exceeded this fall in the theoretical 50% scenario.

Table 2. Comparison of the projected date that ICU capacity is reached, the date ICU demand peaks, the estimated number of ICU beds needed at the peak, and the cumulative COVID-19 deaths at different levels of social distancing. Based on CDC recommendations, in the 50 – 65% scenarios, half of older adults (age 65+) are assumed to practice high levels of social distancing (80%). Projections are showing assuming mask wearing and case isolation remain at current levels.

	Date ICU Capacity Reached*	Date of ICU Peak	ICU Need at Peak**	Cumulative cases through 12/31/2020**	Cumulative deaths through 12/31/2020**
Current trajectory					
Current trajectory (8/17 - 8/25, 65%)	N/A	past	past	402,000	2,400
Three-week average (08/03 - 8/25, 72%)	N/A	past	past	334,000	2,100
Theoretical Intervention scenarios					
Social distancing at 50% [¶]	1/30/21	2/22/21	2,100	1,150,000	4,600
Social distancing at 55% [¶]	N/A	3/29/21	1,300	719,000	3,400
Social distancing at 60% [¶]	N/A	5/27/21	570	497,000	2,700
Social distancing at 65% [¶]	N/A	past	past	396,000	2,300

*ICU capacity for COVID-19 patients is estimated to be 1800 in Colorado, a figure provided by the Colorado Department of Public Health and the Environment.

**Estimates are rounded to three significant figures.

[¶] Intervention are modeled assuming social distancing remains at the current estimated level until 9/7, at which point it changes to the indicated value. These intervention scenarios differ from current trajectory scenarios in that they assume half of individuals 65 and older maintain high levels of social distancing.




Figure 9. Projected daily count of new infections (top), hospital demand (middle), and intensive care (ICU) demand (bottom) in the near-term (left) and long-term (right), assuming social distancing remains at current levels (65%), or hypothetically switches to 65, 60, 55 or 50% on 9/04. Projections are shown assuming mask wearing remains at 90% starting from 7/16, the date of the state-wide mask order. Based on CDC recommendations, in the scenarios half of older adults are assumed to practice high levels of social distancing (80%). Case isolation is assumed to remain at current levels.

Appendix

Code for our model is available on GitHub: <https://github.com/agb85/covid-19>

Model simulations evaluating the potential impact of interventions can be generated using our app: <https://cucovid19.shinyapps.io/colorado/>. This site also includes detailed documentation of our model. The app is updated approximately weekly to reflect our most recent parameter estimates.

Appendix Table A1. Estimated model parameters based on fitting our model output of total hospitalizations to reported hospitalizations in Colorado.

	Range of possible values	Fitted value	Fit using data through
Social distancing*			
Estimated social distancing level over past three weeks, 08/03 - 08/25	30-99%	72%	09/07
Estimated current social distancing level, 08/17 - 08/25	30-99%	65%	09/07
Mask wearing			
Percent of individuals wearing masks, 4/4 to 4/27		50%	Assumed¶
Percent of individuals wearing masks, 4/27 to present		70%	Assumed¶
Percent of individuals wearing masks, 7/16 to present		90%	Assumed
Case isolation			
Decrease in infectious - symptomatic contact rate due to self-isolation by symptomatic after March 5 (dividing by 0.57 gives proportion that self-isolate) **	0.1 - 0.8††	0.44	06/24
Case detection			
Proportion of cases detected over the last 14 days	0.0 – 1.0	42%	08/10
Transmission parameters			
The rate of infection (beta)	0.2 - 0.6††	0.48	06/24
Ratio of infectiousness for symptomatic vs. asymptomatic individuals (lambda)	1.0 - 4.0††	1.39	06/24

*The social distancing parameter estimates the percent decrease in effective contacts between susceptible and infectious individuals. This parameter accounts for social distancing policies intended to avoid contact altogether (e.g., through workplace and school closures) as well as policies and individual behaviors to reduce potential contact with the virus (e.g., maintaining at least 6 feet of distance between people outside of one's household, and handwashing).

† Social distancing estimated weekly and averaged over time period of interest.

¶ Given the difficulty in disentangling the effect of mask wearing from social distancing in decreasing transmission, we cannot fit this parameter at this time. Survey data suggests levels of mask wearing in June were approximately 70% [1]. Recent data from local public health agencies suggest mask wearing is above 90% in stores.

**Self-isolation by symptomatic cases is assumed to occur 1 day after the onset of infectiousness and decrease the 67% of contacts that typically occur outside of the home. This parameter jointly accounts for the percent of symptomatic individuals that self-isolate and the imperfect decline in contacts. Dividing the value in the table by 0.57 gives the proportion of symptomatic individuals that self-isolate.

†† The range of potential parameter values for case isolation [2] and the rate of infectiousness for symptomatic vs. asymptomatic individuals [3, 4] are based on the literature, and for the rate of infection, were obtained from the MIDAS Online COVID-19 compilation of parameter estimates [5].

References

1. Albright E, Banacos N, Birkland T, Crow D, DeLeo R, Dickinson KL, et al. Covid-19 Technical Report Wave One. published online <https://www.riskandsocialpolicy.org/our-work> June 22. 2020.
2. Ferguson N, Laydon D, Nedjati-Gilani G, Imai N, Ainslie K, Baguelin M, et al. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. 2020. Available: <https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-COVID19-NPI-modelling-16-03-2020.pdf>.
3. Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science*. 2020;368(6490):489-93. Epub 2020/03/18. doi: 10.1126/science.abb3221. PubMed PMID: 32179701; PubMed Central PMCID: PMC7164387.
4. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *The New England journal of medicine*. 2020;382(12):1177-9. Epub 2020/02/20. doi: 10.1056/NEJMc2001737. PubMed PMID: 32074444; PubMed Central PMCID: PMC7121626.
5. MIDAS. MIDAS Online COVID-19 Portal 2020. Available from: https://github.com/midas-network/COVID-19/tree/master/parameter_estimates/2019_novel_coronavirus.